



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
6-Methoxy-2-Naphthylacetic acid <i>aka Nabumetone metabolite</i>	Human Plasma	0.5-100 µg/mL	HPLC/FL	0.05 mL	EDTA	-20	Richmond	Inactive	LC184
3-Hydroxymandelic Acid	Human Urine	100-5000 ng/mL	LC/MS/MS	0.05 mL	None	-70	Madison	Active	P957.00
Acetaminophen Acetaminophen (Acetaminophen sulfate) (p-Acetamidophenyl-β-D-glucuronide)	Human Plasma	0.025-10 µg/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Richmond	Active	LC12.1
	Human Urine	0.5-250 µg/mL	HPLC/UV	0.2 mL	None	-20	Richmond	Inactive	LC12.3
	Human Plasma	0.2-50 µg/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Madison	Active	P089.03
	Human Plasma	0.2-50 µg/mL	HPLC/UV	0.1 mL	EDTA	-20	Madison	Active	P829.00
	Dog Plasma	0.2-100 µg/mL	HPLC/UV	0.3 mL	Sodium Heparin	-20	Madison	Active	P363.00
	Human Urine	0.2-200 µg/mL 0.5-500 µg/mL 1-1,000 µg/mL	LC/MS/MS	0.025 mL	None	-20	Madison	Active	P851.00
	Human Plasma	0.1-50 µg/mL 0.1-50 µg/mL 0.5-100 µg/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Madison	Active	P852.00
Acyclovir	Human Plasma	5-1,000 ng/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Richmond	Inactive	LC189
Adiponectin	Human Serum	185-23,695 ng/mL	ELISA	0.01 mL	None	-70	Richmond	Active	ICD287
Albendazole (Sulfoxide & Sulfone Metabolites)	Bovine Plasma	0.05-5 µg/mL 0.05-5 µg/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Madison	Active	P562.02

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Albuterol	Human Plasma	0.20-20 ng/mL	LC/MS/MS	1.0 mL	Sodium Heparin	-20	Madison	Active	P478.01
	Human Plasma	2-4,000 pg/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Madison	Active	P951.00
	Human Urine	20-2,000 ng/mL	LC/MS/MS	0.1 mL	None	-20	Madison	Active	P465.01
	Dog Plasma	0.2-20 ng/mL	LC/MS/MS	1.0 mL	Sodium Heparin	-20	Madison	Active	P464.01
	Rat Plasma	0.4-400 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Madison	Inactive	P463.00
Alprazolam	Human Plasma	0.25-40 ng/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Richmond	Active	LCMS225
Alprazolam (α -Hydroxyalprazolam) (4-Hydroxyalprazolam)	Human Plasma	0.2-40 ng/mL 0.2-40 ng/mL 0.2-40 ng/mL	LC/MS/MS	0.3 mL	Sodium Heparin	-20	Madison	Active	P335.06
Ambrisentan (4-Hydroxymethyl Ambrisentan)	Human Plasma	1-400 ng/mL 0.25 -100 ng/mL	LC/MS/MS	0.1 mL	Sodium Citrate	-20	Richmond	Active	LCMSC498
	Human Plasma	1-500 ng/mL 0.25-125 ng/mL	LC/MS/MS	0.1 mL	K₂EDTA	-20	Richmond	Active	LCMSC498.1
Amitriptyline (Nortriptyline)	Human Plasma	0.5-200 ng/mL 0.5-200 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Madison	Active	P558.01
	Rabbit Plasma	1-500 ng/mL 1-500 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P647.00
Amitriptyline	Human Plasma	0.5-200 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Richmond	Active	LCMS301

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Amlodipine	Human Plasma	0.1-50 ng/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-20	Richmond	Active	LCMS151
Amoxicillin	Human Serum	50-25,000 ng/ml	LC/MS/MS	0.1 mL	None	-70	Richmond	Active	LCMS141
	Human Plasma	50-25,000 ng/ml	LC/MS/MS	0.1 mL	K ₃ EDTA	-70	Richmond	Active	LCMS141.2
Atazanavir & Ritonavir	Human Plasma	10-10,000 ng/mL	LC/MS/MS	0.05 mL	K₃EDTA	-20	Madison	Active	P1012
Atenolol	Human Whole Blood	10-1,000 ng/mL	HPLC/FL	1.0 mL	Sodium Heparin	-20	Madison	Inactive	P161.01
Aripiprazole (Dehydroaripiprazole)	Human Plasma	0.5-250 ng/mL 0.1-50 ng/mL	LC/MS/MS	0.25 mL	K ₂ EDTA	-20	Richmond	Active	LCMS 386
Atovaquone	Human Plasma	0.025-10 µg/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Richmond	Active	LCMS360
Atorvastatin (Ortho-hydroxy atorvastatin) (Para-hydroxy atorvastatin)	Human Plasma	0.1-75 ng/mL 0.1-75 ng/mL 0.1-75 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-70	Richmond	Active	LCMS120
	Human Plasma	0.1-75 ng/mL 0.1-75 ng/mL 0.1-75 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-70	Madison	Active	P913.00
	Rat Plasma	0.1-75 ng/mL 0.1-75 ng/mL 0.1-75 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-70	Richmond	Active	LCMS120.3
Avastin aka Bevacizumab	Human Serum	10-2,560 ng/mL	MSD-ECL	0.025 mL	None	-70	Richmond	Active	ICD343
Azithromycin	Human Plasma	2.5-1,000 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Richmond	Inactive	LCMS168

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Azidothymidine (AZT)	Human Plasma	50-25,000 ng/mL	HPLC/UV	0.25 mL	Sodium Heparin	-20	Richmond	Inactive	LC96
	Human Urine	0.2-25 µg/mL	HPLC/UV	1.0 mL	None	-20	Richmond	Inactive	LC96.1
Beclomethasone (Beclomethasone-17-propionate) (Beclomethasone dipropionate)	Human Plasma	1-250 pg/mL 1-250 pg/mL 1-250 pg/mL	LC/MS/MS	1.0 mL	Potassium Oxalate/Sodium Fluoride	-70	Richmond	Active	LCMSC333
Benazepril (Benazeprilat)	Human Plasma	10-1,000 ng/mL 10-1,000 ng/mL	LC/MS/MS	0.05 mL	Na ₂ EDTA	-20	Richmond	Active	LCMS107
Bicalutamide	Human Plasma	10-1,250 ng/mL	LC/MS/MS	0.05 mL	K ₃ EDTA	-20	Richmond	Active	LCMS318
Bisoprolol	Human Plasma	0.5-200 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Richmond	Active	LCMSB501
Bortezomib	Human Plasma	0.5-500 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-70	Richmond	Active	LCMS509
Brompheniramine	Human Plasma	0.05-20 ng/mL	LC/MS/MS	0.1 mL	Lithium Heparin	-20	Richmond	Active	LCMSC436
Budesonide	Human Plasma	10-2,500 pg/mL	LC/MS/MS	0.25 mL	K ₃ EDTA	-20	Madison	Active	P429.00
Buprenorphine (Norbuprenorphine) & Naloxone	Human Plasma	50-10,000 pg/mL 50-10,000 pg/mL 5-500 pg/mL	LC/MS/MS	0.5 mL	K ₃ EDTA	-20	Madison	Active	P942.01
Bupropion (Hydroxybupropion) (Erythrohydroxybupropion) (Threoxyhydroxybupropion)	Human Plasma	0.5-200 ng/mL 1-400 ng/mL 1-400 ng/mL 1-400 ng/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-70	Madison	Active	P986.00

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Butalbital	Human Plasma	0.05-25 µg/mL	HPLC/UV	0.25 mL	Sodium Heparin	-20	Richmond	Active	LC196
	Human Plasma	0.1-10 µg/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Madison	Inactive	P301.00
Caffeine (1,7-Dimethylxanthine)	Human Plasma	20-20,000 ng/mL 20-20,000 ng/mL	LC/MS/MS	0.15 mL	Sodium Heparin	-20	Richmond	Active	LCMSC207
	Human Urine	1-50 µg/mL 1-50 µg/mL	HPLC/UV	0.1 mL	None	-20	Madison	Active	P590.02
	Human Plasma	25-25,000 ng/mL 25-25,000 ng/mL	HPLC/UV	0.25 mL	Sodium Heparin	-20	Madison	Active	P589.02
	Human Plasma	25-25,000 ng/mL 25-25,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Madison	Active	P734.03
	Human Urine	50-50,000 ng/mL 50-50,000 ng/mL	LC/MS/MS	0.025 mL	None	-20	Madison	Active	P739.03
Candesartan	Human Plasma	0.5-400 ng/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Richmond	Active	LCMSC451
	Human Urine	10-5,000 ng/mL	LC/MS/MS	0.05 mL	None	-20	Richmond	Active	LCMSC451.1
Carbidopa & Levodopa	Human Plasma	0.5-250 ng/mL 5-2,500 ng/mL	LC/MS/MS	0.2 mL	K ₃ EDTA	-70	Madison	Active	P838.00
Carbidopa (3-O-MethylDopa) & Levodopa	Human Plasma	0.5-250 ng/mL 25-25,000 ng/mL 10-5,000 ng/mL	LC/MS/MS	0.2 mL	K ₃ EDTA	-70	Madison	Active	P959.00

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Carbamazepine (Carbamazepine 10,11 Epoxide)	Human Plasma	0.05-20 µg/mL 0.05-20 µg/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Madison	Active	P239.03
	Human Plasma	0.05-50 µg/mL 0.05-50 µg/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Madison	Active	P626.00
Carbamazepine, Free (unbound) (Carbamazepine 10,11 Epoxide) (Carbamazepine 10,11 Epoxide) (Monohydroxycarbazepine)	Human Plasma Ultrafiltrate	0.1-20 µg/mL 0.1-20 µg/mL	HPLC/UV	0.25 mL	Sodium Heparin	-20	Madison	Inactive	P238.02
	Human Urine	0.1-20 µg/mL	HPLC/UV	0.25 mL	None	-20	Madison	Inactive	P257.01
	Human Plasma	1-50 µg/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-70	Madison	Active	P886.00
Carvedilol	Human Plasma	0.1-50 ng/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-20	Madison	Active	P789.01
Cefaclor	Human Plasma	0.1-25 µg/mL	HPLC/UV	0.5 mL	Sodium Heparin	-70	Richmond	Active	LC305
Cefadroxil	Human Serum	0.25-50 µg/mL	HPLC/UV	0.10 mL	None	-70	Richmond	Inactive	LC52
	Human Urine	5-1,000 µg/mL	HPLC/UV	0.05 mL	None	-20	Richmond	Inactive	LC52.1
Celecoxib	Human Plasma	5-2,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Richmond	Active	LCMSD417
Celiprolol	Human Plasma	10-2,000 ng/mL	HPLC/UV	1.0 mL	Sodium Heparin	-20	Richmond	Inactive	LC30
Cephalexin	Human Plasma	0.1-100 µg/mL	LC/MS/MS	0.05 mL	K ₃ EDTA	-20	Richmond	Active	LCMSB382
Cetirizine	Human Plasma	0.5-500 ng/mL	LC/MS/MS	0.05 mL	K ₃ EDTA	-20	Richmond	Active	LCMSB169

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Chlorpheniramine	Human Plasma	0.25-50 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Richmond	Active	LCMS214
Chlorthalidone	Human Plasma	2-1,000 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Richmond	Active	LCMS389.2
	Human Urine	10-5,000 ng/mL	LC/MS/MS	0.1 mL	None	-20	Richmond	Active	LCMS389.1
Chlorzoxazone (6-Hydroxychlorzoxazone)	Human Plasma	5-5,000 ng/mL 5-5,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Richmond	Active	LCMS209
Cimetidine	Human Urine	5.0-5,000 µg/mL	HPLC/UV	0.05 mL	None	-20	Richmond	Active	LC358.1
	Human Plasma	0.01-10 µg/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Richmond	Active	LC358
Ciprofloxacin	Human Plasma	25-10,000 ng/mL	LC/MS/MS	0.05 mL	K ₃ EDTA	-20	Richmond	Active	LCMS176
Citalopram	Human Plasma	1-500 ng/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-20	Richmond	Active	LCMS332
Citalopram (Desmethylcitalopram)	Human Plasma	1-500 ng/mL 0.2-100 ng/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-20	Madison	Active	P787.00
Clarithromycin (14-Hydroxyclearithromycin)	Human Plasma	20-10,000 ng/mL 5-2,500 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Richmond	Active	LCMSB179
Clavulanic Acid	Feline Plasma	0.1-10 µg/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-80	Madison	Active	P997.00
	Canine Plasma	0.1-10 µg/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-70	Madison	Active	P990.00
Clopidogrel (Clopidogrel Acid)	Human Plasma	0.2-200 ng/mL 10-10,000 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Richmond	Active	LCMS271
Clorsulon	Bovine Plasma	0.1-10 ng/mL	LC/MS/MS	1.0 mL	Sodium Heparin	-20	Madison	Active	P717.00

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Cocktail Assay, "Epilepsy" Carbamazepine Lamotrigine Phenobarbital Phenytoin Topiramate	Human Plasma	0.1-50 µg/mL 0.1-20 µg/mL 0.1-50 µg/mL 0.1-30 µg/mL 0.1-50 µg/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P837.00
Cocktail Assay, "Antiviral" Darunavir Lopinavir Raltegravir Ritonavir	Human Plasma	10-10,000 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Madison	Active	P1002.01
Codeine	Human Plasma	0.1-100 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Madison	Active	P753.01
Codeine & Morphine (Morphine-3-β-glucuronide) (Morphine-6-β-glucuronide)	Human Plasma	1-100 ng/mL 0.2-20 ng/mL 2-200 ng/mL 0.5-50 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Richmond	Active	LCMSC396
Codeine & Morphine & Guafenesin	Human Plasma	0.1-100 ng/mL 0.2-20 ng/mL 2-2,000 ng/mL	LC/MS/MS	0.3 mL	Sodium Heparin	-20	Madison	Active	P919.02
Cortisol (6β-Hydroxycortisol)	Human Urine	1-200 ng/mL 6-1,200 ng/mL	LC/MS/MS	0.1 mL	None	-20	Richmond	Active	LCMSC355
Cortisol <i>aka hydrocortisone</i> Cortisone	Human Urine	1-250 ng/mL 1-250 ng/mL	LC/MS/MS	0.05 mL	None Requires CHAPS	-70	Richmond	Active	LCMSC472

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Cortisol <i>aka hydrocortisone</i> Cortisone	Human Serum	1-500 ng/mL 0.5-250 ng/mL	LC/MS/MS	0.05 mL	None	-70	Richmond	Active	LCMSD472.1
5- α -Tetrahydrocortisol (5- α -THF) 5- β -Tetrahydrocortisol (5- β -THF) Tetrahydrocortisone (THE)	Human Urine	20-5,000 ng/mL 20-5,000 ng/mL 20-5,000 ng/mL	LC/MS/MS	0.05 mL	None Requires CHAPS	-70	Richmond	Active	LCMSC471
Cyclophosphamide	Human Plasma	0.25–50 μ g/mL	HPLC/UV	0.25 mL	Sodium Heparin	-20	Richmond	Active	LC232
D4T	Human Plasma	0.025-25 μ g/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Richmond	Inactive	LC211
Danazol	Human Plasma	2-200 ng/mL	HPLC/UV	1.0 mL	Sodium Heparin	-20	Richmond	Inactive	LC18
Darunavir See "Cocktail Assay Antiviral"	Human Plasma	10-10,000 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Madison	Active	P1002.00
Daunorubicin (Daunorubicinol)	Human Plasma	5-200 ng/mL 5-200 ng/mL	HPLC/FL	0.5 mL	Sodium Heparin	-20	Richmond	Inactive	LC301
Desipramine (2-Hydroxydesipramine)	Human Plasma	0.25-100 ng/mL 0.25-100 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Richmond	Active	LCMS103
	Human Plasma	0.25-100 ng/mL 0.25-100 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P644.01
Dexamethasone	Human Plasma	0.5-500 ng/mL	LC/MS/MS	0.2 mL	K ₂ EDTA	-20	Richmond	Active	LCMS328
Dexlansoprazole	Human Plasma	5-5,000 ng/mL	LC/MS/MS	0.05 mL	K₂EDTA	-20	Madison	Active	P1026
Dextromethorphan	Human Plasma	10 -5,000 pg/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Madison	Active	P732.01

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Dextromethorphan (Dextrophan)	Dog Plasma	2.5-500 ng/mL 2.5-500 ng/mL	HPLC/FL	0.2 mL	Sodium Heparin	-20	Madison	Active	P423.00
	Rat Plasma	5-1,000 ng/mL 5-1,000 ng/mL	HPLC/FL	0.1 mL	Sodium Heparin	-20	Madison	Inactive	P422.00
	Human Plasma	0.05- 50 ng/mL 0.8-800 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P668.05
Dextromethorphan, Total** & Unconjugated (Dextrophan) (3-methoxymorphinan) (3-hydroxymorphinan)	Human Urine	0.001-1.0 µg/mL 0.020-20 µg/mL 0.001-1.00 µg/mL 0.01-10.0 µg/mL	LC/MS/MS	0.05 mL	None	-20	Madison	Active	P600.07
Dextromethorphan & Pseudoephedrine	Human Plasma	10-5,000 pg/mL 0.8-400 ng/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Madison	Active	P617.02
Diazepam (N-Desmethyldiazepam)	Human Plasma	2-1,000 ng/mL 1-500 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Richmond	Active	LCMS 466
Diclofenac	Human Plasma	0.1-100 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Richmond	Active	LCMSC368
Diclofenac (4'-hydroxydiclofenac)	Human Plasma	10-2,000 ng/mL 2.5-500 ng/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-20	Richmond	Active	LCMSB368.1
Digoxin	Human Plasma	0.15-8 ng/mL	RIA	0.2 mL	Sodium Heparin	-70	Richmond	Active	ICD11.1
	Human Serum	0.15-8 ng/mL	RIA	0.1 mL	None	-70	Richmond	Active	ICD11.3
	Human Urine	1-40 ng/mL	RIA	0.1 mL	None	-70	Richmond	Active	ICD11.2

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Digoxin	Human Plasma	0.1-4 ng/mL	RIA	0.4 mL	EDTA	-70	Richmond	Active	ICD11.5
	Human Plasma	0.1-50 ng/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Richmond	Active	LCMSC455.1
	Human Plasma	0.05-50 ng/mL	LC/MS/MS	0.1 mL	K₂EDTA	-20	Richmond	Active	LCMSD455.2
Diltiazem (Desacetyldiltiazem) (Desmethyldiltiazem)	Human Plasma	0.1-200 ng/mL 0.1-200 ng/mL 0.1-200 ng/mL	LC/MS/MS	0.4 mL	Sodium Heparin	-20	Madison	Active	P672.03
Docetaxel	Human Plasma	10-5,000 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Richmond	Active	LCMSB216
Donepezil	Human Plasma	0.2-50 ng/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-20	Richmond	Active	LCMSB203
	Human Plasma	0.2-100 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P983.00
Doxazosin	Human Plasma	0.05-50 ng/mL	HPLC/FL	1.0 mL	Sodium Heparin	-20	Richmond	Inactive	LC266
Doxepin (Desmethyldoxepin)	Human Plasma	0.5-250 ng/mL 0.5-250 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P815.01
Doxorubicin	Human Plasma	1-200 ng/mL	HPLC/FL	0.5 mL	Sodium Heparin	-70	Richmond	Inactive	LC252
Duloxetine	Human Plasma	0.2-200 ng/mL	LC/MS/MS	0.5 mL	K ₂ EDTA	-20	Madison	Active	P882.00
Efavirenz	Human Plasma	0.10-10 µg/mL	LC/MS/MS	0.1 mL	K₂EDTA	-20	Richmond	Active	LCMSB517
	Human Whole Blood	0.10-25 µg/mL	DBS by LC/MS/MS	0.1 to 0.3 mL	K₂EDTA	Room Temp	Richmond	Active	LCMS517.1
Emtricitabine & Tenofovir	Human Plasma	20-4,000 ng/mL 5-1,000 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Richmond	Active	LCMSC 468

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Emtricitabine & Tenofovir	Human Plasma	5-1,000 ng/mL 20-4,000 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Madison	Active	P947.01
Enalapril (Enalaprilat)	Human Plasma	0.25-100 ng/mL 0.25-100 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Madison	Active	P435.00
	Human Plasma	0.5-250 ng/mL 0.5-250 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Richmond	Inactive	LCMS31
	Dog Plasma	0.25-400 ng/mL 0.25-400 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Madison	Active	P532.01
Epinephrine	Human Plasma	0.05-8 ng/mL	HPLC/FL	1.0 mL	Sodium Metabisulfite	-20	Madison	Active	P981.02
Erlotinib	Human Plasma	10-10,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Richmond	Active	LCMSC390
	Human Plasma	10-10,000 ng/mL	LC/MS/MS	0.025 mL	Sodium Heparin	-20	Madison	Active	P1008.01
	Human Plasma	5-5,000 ng/mL	LC/MS/MS	0.025 mL	Sodium Heparin	-20	Madison	Active	P1016
Erythromycin	Human Plasma	0.2-20 µg/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Richmond	Active	LCMS463
Estazolam	Human Plasma	0.1-30 ng/mL	LC/MS/MS	1.0 mL	Sodium Heparin	-20	Madison	Inactive	P336.00
β-Estradiol & Equilin & Estrone	Human Plasma	5.0-500 pg/mL 5.0-500 pg/mL 5.0-500 pg/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-70	Richmond	Active	LCMSC233

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
β-Estradiol & Equilin & Estrone	Human Plasma	1.0-100 pg/mL 5.0-500 pg/mL 5.0-500 pg/mL	LC/MS/MS	0.5 mL	K ₃ EDTA	-70	Richmond	Active	LCMSC350
	Human Serum	1.0-100 pg/mL 5.0-500 pg/mL 5.0-500 pg/mL	LC/MS/MS	0.5 mL	None	-70	Richmond	Active	LCMSC350.1
β-Estradiol (Estrone)	Human Serum	2.5 – 250 pg/mL 5.0 – 500 pg/mL	LC/MS/MS	0.5 mL	None	-20	Richmond	Inactive	LCMSC248.1
	Human Plasma	2.5 – 250 pg/mL 5 – 500 pg/mL	LC/MS/MS	0.5 mL	K ₃ EDTA	-70	Richmond	Active	LCMSC248
β-Estradiol, Total**	Human Serum	0.025-10 ng/mL	LC/MS/MS	0.1 mL	None	-70	Richmond	Active	LCMS295
Estrone Sulfate	Human Serum	50-5,000 pg/mL	LC/MS/MS	0.5 mL	None	-70	Richmond	Active	LCMS27.1
Estrone & Equilin, Total**	Human Plasma	10-5,000 pg/mL 10-5,000 pg/mL	LC/MS/MS	0.5 mL	K ₃ EDTA	-70	Richmond	Active	LCMSC351
	Human Serum	10-5,000 pg/mL 10-5,000 pg/mL	LC/MS/MS	0.5 mL	None	-70	Richmond	Active	LCMSC351.1
Ethanol	Human Plasma	20-1,000 µg/mL	GC/FID	0.1 mL	Potassium Oxalate/Sodium Fluoride	-20	Richmond	Active	GC89
Ethinyl Estradiol & Norethindrone	Human Plasma	2-500 pg/mL 50-25,000 pg/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Richmond	Active	LCMSC255

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Ethinyl Estradiol & Norgestrel	Human Plasma	2-500 pg/mL 50-25,000 pg/mL	LC/MS/MS	0.5 mL	Potassium Oxalate/Sodium Fluoride	-70	Richmond	Active	LCMSC256
Ethinyl Estradiol Sulfate	Human Plasma	50-10,000 pg/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Richmond	Active	LCMSC398
Etodolac	Human Plasma	0.2-50 µg/mL	HPLC/UV	1.0 mL	Sodium Heparin	-20	Richmond	Active	LC193
Etoposide	Human Plasma	0.02-20 µg/mL	LC/MS/MS	0.05 mL	K₂EDTA	-20	Richmond	Active	LCMS515
Etoricoxib	Human Plasma	5-2500 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Madison	Active	P971.01
Everolimus	Human Whole Blood	0.2-200 ng/mL	LC/MS/MS	0.5 mL	K₂EDTA	-20	Madison	Active	P1022
Ezetimibe	Human Plasma	0.2-100 ng/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Richmond	Active	LCMSC439
Ezetimibe, Total	Human Plasma	0.2-100 ng/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Madison	Active	P945.00
	Human Plasma	1.0-500 ng/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Richmond	Active	LCMSC440
	Human Plasma	1.0-500 ng/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Madison	Active	P944.00
	Human Plasma	1.0-500 ng/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Madison	Active	P944.00
Famotidine	Human Plasma	5-250 ng/mL	HPLC/UV	1.0 mL	Sodium Heparin	-20	Madison	Inactive	P255.01
Fenofibric Acid	Human Plasma	0.05-30 µg/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Richmond	Active	LCMSB235
Fentanyl	Human Plasma	25-10,000 pg/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Richmond	Active	LCMSB186.5
	Human Plasma	5-2,000 pg/mL	LC/MS/MS	0.25 mL	K ₃ EDTA	-20	Richmond	Active	LCMSC186.4
	Human Serum	10-2,000 ng/mL	LC/MS/MS	0.1 mL	None	-20	Madison	Active	P869.01
	Dog Plasma	40-40,000 pg/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Madison	Active	P966.00

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Fentanyl (Norfentanyl)	Mouse Plasma	0.04-40 ng/mL 0.04-40 ng/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Madison	Active	P637.02
	Rat Plasma	0.04-40 ng/mL 0.04-40 ng/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Madison	Active	P823.01
	Rabbit Plasma	0.04-40 ng/mL 0.04-40 ng/mL	LC/MS/MS	0.2 mL	K ₂ EDTA	-20	Madison	Active	P636.01
Fentanyl (Norfentanyl) (DespropionylFentanyl)	Human Plasma	10-500 pg/mL 10-500 pg/mL 10-500 pg/mL	LC/MS/MS	0.5 mL	K ₃ EDTA	-20	Madison	Active	P624.02
Fexofenadine	Human Plasma	0.5-500 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P745.04
	Human Urine	0.05-10 µg/mL	LC/MS/MS	0.02 mL	None	-20	Madison	Active	P746.04
Fluconazole	Human Plasma	20-10,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Madison	Active	P791.00
Flunixin	Equine Plasma	10-5,000 ng/mL	LC/MS/MS	0.025 mL	Sodium Heparin	-70	Madison	Active	P970.00
Fluoxetine (Norfluoxetine)	Human Plasma	0.25-100 ng/mL 0.25-100 ng/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-20	Richmond	Active	LCMS52
Fluticasone Propionate	Human Plasma	2.5-500 pg/mL	LC/MS/MS	1.0 mL	Potassium Oxalate/Sodium Fluoride	-70	Richmond	Active	LCMSC173
	Human Plasma	1.0-500 pg/mL	LC/MS/MS	1.0 mL	Potassium Oxalate/Sodium Fluoride	-20	Madison	Active	P817.02
Fluticasone & Salmeterol	Human Plasma	1-200 pg/mL 1-200 pg/mL	LC/MS/MS	1.0 mL	Potassium Oxalate/Sodium Fluoride	-20	Madison	Active	P988

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Fluvoxamine	Human Plasma	0.5-500 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P658.01
Formoterol	Human Plasma	0.5-250 pg/mL	LC/MS/MS	1.0 mL	Lithium Heparin with Eserine hemisulfate	-20	Madison	Active	P860.02
	Human Plasma	0.25-125 pg/mL	LC/MS/MS	1.0 mL	Lithium Heparin with Eserine hemisulfate	-20	Madison	Active	P996.00
	Human Urine	6.0-1,200 pg/mL	LC/MS/MS	0.1 mL	None	-20	Madison	Active	P883.01
Formoterol Enantiomers (R,R and S,S)	Human Urine	6.0-1,200 pg/mL each	LC/MS/MS	0.5 mL	None	-20	Madison	Active	P868.00
Furosemide	Human Urine	0.05-15 µg/mL	HPLC/FL	1.0 mL	None	-20	Madison	Active	P376.00
	Human Plasma	0.02-10 µg/mL	HPLC/FL	0.5 mL	Sodium Heparin	-20	Madison	Active	P372.01
	Human Plasma	0.02-10 µg/mL	HPLC/FL	0.2 mL	Sodium Heparin	-20	Richmond	Active	LC354
	Human Urine	0.2-50 µg/mL	LC/MS/MS	0.2 mL	None	-20	Richmond	Active	LCMS317
Gabapentin	Human Plasma	0.05- 10 µg/mL	LC/MS/MS	0.2 mL	K ₃ EDTA	-20	Richmond	Active	LCMS359
Galanthamine	Human Plasma	0.1-100 ng/mL	LC/MS/MS	0.25 mL	K ₃ EDTA	-20	Richmond	Active	LCMSB312
Gatifloxacin	Human Plasma	0.025-10 µg/mL	LC/MS/MS	0.05 mL	K ₃ EDTA	-20	Richmond	Active	LCMSB285
Gemfibrozil	Human Plasma	0.05-25 µg/mL	HPLC/FL	0.5 mL	Sodium Heparin	-20	Richmond	Inactive	LC56
Gemfibrozil	Human Plasma	0.05-50 µg/mL	HPLC/UV	0.5 mL	Sodium Heparin and EDTA	-20	Madison	Inactive	P115.00

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Glimepiride	Human Plasma	2-1,000 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Richmond	Active	LCMS266
Glipizide	Human Plasma	1-500 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Richmond	Active	LCMS137
Glyburide	Human Plasma	1-500 ng/mL	LC/MS/MS	0.25 mL	K ₃ EDTA	-20	Richmond	Active	LCMS159
Glyburide <i>Lower range for pediatric studies</i>	Human Plasma	0.10-50 ng/mL	LC/MS/MS	0.25 mL	K ₃ EDTA	-20	Richmond	Active	LCMS159.1
Guaifenesin	Human Plasma	2-2,000 ng/mL	LC/MS/MS	0.075 mL	Sodium Heparin	-20	Madison	Active	P973.00
Guaifenesin & Codeine & Morphine	Human Plasma	2-2,000 ng/mL 0.1-100 ng/mL 0.2-20 ng/mL	LC/MS/MS	0.3 mL	Sodium Heparin	-20	Madison	Active	P919.02
Herceptin <i>aka Trastuzumab</i>	Human Serum	10-200 µg/mL	ELISA	0.2 mL	None	-70	Richmond	Active	ICD178
	Human Plasma	10-100 µg/mL	ELISA	0.1 mL	EDTA	-20	Richmond	Active	ICD178.1
Hydrochlorothiazide	Human Plasma	2-200 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Richmond	Active	LCMSB205.4
	Human Plasma	2-200 ng/mL	LC/MS/MS	0.25 mL	K ₃ EDTA	-20	Richmond	Active	LCMSB205
	Human Urine	0.1-25 µg/mL	LC/MS/MS	0.05 mL	None	-20	Richmond	Active	LCMSB205.3
Hydrocodone (Hydromorphone)	Human Plasma	0.1-100 ng/mL 0.05-50 ng/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Richmond	Active	LCMSC492
Hydrocodone (Hydromorphone)	Human Plasma	0.1-100 ng/mL 0.05-50 ng/mL	LC/MS/MS	0.5 mL	K ₃ EDTA	-20	Madison	Inactive	P623.00

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
	Hamster Plasma	1-400 ng/mL 1-400 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Madison	Inactive	P659.00
Hydrocodone & Ibuprofen	Human Plasma	0.25-50 ng/mL 0.25-50 ug/mL	LC/MS/MS	0.2 mL	K ₃ EDTA	-20	Madison	Active	P774.01
Ibuprofen	Human Plasma	0.2-50 µg/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Richmond	Active	LCMSB409
	Human Plasma	0.2-50 µg/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Madison	Inactive	P046.04
IFN Gamma	Human Serum	31.3-1,000 pg/mL	ELISA	0.2 mL	None	-70	Richmond	Active	ICD278
IL-1β	Human Serum	7.82-250 pg/mL	ELISA	0.4 mL	None	-20	Richmond	Active	ICD151
IL-2	Human Serum	62.5-2,000 pg/mL	ELISA	0.2 mL	None	-20	Richmond	Active	ICD150
IL-4	Human Serum	31.3-2,000 pg/mL	ELISA	0.1 mL	None	-70	Richmond	Active	ICD277
IL-6	Human Serum	5-320 pg/mL	ELISA	0.2 mL	None	-20	Richmond	Active	ICD141
IL-10	Human Serum	10- 500 pg/mL	ELISA	0.3 mL	None	-20	Richmond	Active	ICD142
Imantinib	Human Plasma	10-10,000 ng/mL	LC/MS/MS	0.05 mL	K₂EDTA	-20	Madison	Active	P1010
Irbesartan	Human Plasma	2.5-2,500 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Richmond	Active	LCMSB232
Isopropamide	Human Plasma	0.05-10 ng/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Madison	Inactive	P409.01
Itraconazole (2-Hydroxyitraconazole)	Human Plasma	0.5-100 ng/mL 0.5-100 ng/mL	HPLC/FL	1.0 mL	Sodium Heparin	-20	Madison	Active	P476.01

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Ivermectin B1a	Equine Plasma	0.5-50 ng/mL	HPLC/FL	0.5 mL	Sodium Heparin	-20	Madison	Active	P575.01
	Equine Serum	0.50-50 ng/mL 0.84-1.68 ng/mL	HPLC/FL	2.0 mL	None	-20	Madison	Active	P389.00
Ivermectin B1b	Equine Plasma	0.10-10 ng/mL	HPLC/FL	0.5 mL	Sodium Heparin	-20	Madison	Active	P621.00
Ivermectin B1a	Swine Plasma	0.50-100 ng/mL	HPLC/FL	0.5 mL	Sodium Heparin	-20	Madison	Active	P646.00
	Bovine Plasma	0.50-100 ng/mL	HPLC/FL	0.5 mL	Sodium Heparin	-20	Madison	Active	P479.01
	Dog Plasma	0.50-50 ng/mL	HPLC/FL	0.5 mL	Sodium Heparin	-20	Madison	Active	P457.02
Ivermectin B1a	Dog Plasma	0.10-10 ng/mL	HPLC/FL	0.5 mL	Sodium Heparin	-20	Madison	Active	P715.01
Ketamine (Norketamine)	Human Serum	5-5,000 ng/mL 2.5-2,500 ng/mL	LC/MS/MS	0.2 mL	None	-20	Madison	Active	P627.00
	Rabbit Plasma	10-10,000 ng/mL 5-5,000 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P648.00
	Human Plasma	0.5-500 ng/mL 0.5-500 ng/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-20	Madison	Active	P663.02
S-Ketamine and R-Ketamine S-Norketamine and R-Norketamine	Feline Plasma	20-8,000 ng/mL 10-4,000 ng/mL	LC/MS/MS	50 uL	Sodium Heparin	-20	Madison	Active	P703.01
Ketoconazole	Human Plasma	0.05-5 µg/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Richmond	Active	LCMSB202
Ketoprofen	Human Plasma	1-500 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Richmond	Active	LCMSB162

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Ketoprofen, and Total**	Human Plasma	0.1-50 ng/mL	LC/MS/MS	0.4 mL	Sodium Heparin	-20	Madison	Active	P747.02
	Mini Pig	0.1-50 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P821.00
	Human Plasma	1.0-1,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Madison	Active	P767.00
	Human Urine	5-5,000 ng/mL	LC/MS/MS	0.025 mL	None	-20	Madison	Active	P768.01
Ketoprofen Enantiomers	Human Plasma	0.5– 250 ng/mL 0.5-250 ng/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-20	Madison	Active	P782.00
Ketorolac	Human Plasma	0.025-25 µg/mL	HPLC/UV	0.25 mL	Sodium Heparin	-20	Richmond	Active	LC207
Lamivudine (3TC)	Human Plasma	10-4000 ng/mL	HPLC/UV	0.5 mL	EDTA	-20	Madison	Active	P560.00
	Human Plasma	10-5,000 ng/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Richmond	Active	LC269
Lamotrigine See Cocktail Assay "Epilepsy"	Human Plasma	0.1-20 µg/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-70	Madison	Active	P837.00
Lansoprazole	Human Plasma	2-2,000 ng/mL	LC/MS/MS	0.05mL	K ₃ EDTA	-20	Richmond	Active	LCMS178
	Human Plasma	2-2,000 ng/mL	LC/MS/MS	0.05mL	K ₃ EDTA	-20	Madison	Active	P896.02
Leucovorin Diasteromers (6S-5-Methyltetrahydrofolic Acid Diastereomers)	Human Serum	0.5-250 µg/mL 0.1-50 µg/mL	LC/MS/MS	0.05 mL	None	-70	Richmond	Active	LCMSC387
Leuprolide	Human Plasma	0.025-25 ng/mL	LC/MS/MS	0.5 mL	K ₃ EDTA	-20	Richmond	Active	LCMSC245
Levetiracetam	Human Plasma	0.05-50 µg/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Madison	Active	P811.03

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Levodopa & Carbidopa	Human Plasma	5-2,500 ng/mL 0.5-250 ng/mL	LC/MS/MS	0.2 mL	K ₃ EDTA	-70	Madison	Active	P838.00
Levodopa & Carbidopa (3-O-MethylDopa)	Human Plasma	10-5,000 ng/mL 0.5-250 ng/mL 25-25,000 ng/mL	LC/MS/MS	0.2 mL	K ₃ EDTA	-70	Madison	Active	P959.00
Levofloxacin	Human Serum	25-15,000 ng/mL	HPLC/UV	0.5 mL	None	-20	Richmond	Active	LC287
Lidocaine	Human Plasma	2-500 ng/mL	HPLC/UV	0.25 mL	Sodium Heparin	-20	Madison	Active	P654.00
	Pig Plasma	2-500 ng/mL	HPLC/UV	0.25 mL	Sodium Heparin	-20	Madison	Active	P634.00
	Rat Plasma	1-1,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Madison	Active	P667.00
	Human Plasma	0.2-200 ng/mL	LC/MS/MS	0.05 mL	K ₃ EDTA	-20	Madison	Active	P871.00
Lisinopril	Human Plasma	0.5-400 ng/mL	LC/MS/MS	0.3 mL	Sodium Heparin	-20	Madison	Active	P698.01
	Human Urine	25-25,000 ng/mL	LC/MS/MS	0.025 mL	None	-20	Madison	Active	P694.00
	Rat Plasma	1.0-1,000 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P673.00
Loperamide	Human Plasma	10-10,000 pg/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Madison	Active	P557.02
	Human Plasma	10-10,000 pg/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Richmond	Active	LCMSC310
Lopinavir See Cocktail Assay "Antiviral"	Human Plasma	10-10,000 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Madison	Active	P1002.01
Loratadine (Descarboethoxyloratadine) (3-OHDescarboethoxyloratadine)	Human Plasma	0.025-10 ng/ml 0.025-10 ng/ml 0.025-10 ng/ml	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Richmond	Active	LCMS188.1

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Lorazepam	Human Plasma	0.5-50 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Richmond	Active	LCMS142
Lorazepam, Total**	Human Plasma	2.5-500 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Richmond	Active	LCMSB294
Losartan (Acid Metabolite)	Human Plasma	0.5-500 ng/mL 0.5-500 ng/mL	LC/MS/MS	0.15 mL	Sodium Heparin	-20	Richmond	Active	LCMSC208
	Human Urine	5-2,000 ng/mL 2-2,000 ng/mL	LC/MS/MS	0.1 mL	None	-20	Richmond	Inactive	LCMSC208.1
	Human Plasma	0.05-12.5 ng/mL 0.1-25 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-70	Richmond	Inactive	LCMS267
Meclocycline	Human Plasma	25-10,000 ng/mL	HPLC/UV	0.2 mL	K ₂ EDTA	-20	Richmond	Active	LC337
Medroxyprogesterone Acetate	Human Plasma	0.01-10 ng/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Richmond	Inactive	LCMS99
	Human Serum	0.02-5 ng/mL	LC/MS/MS	0.25 mL	None	-20	Richmond	Active	LCMSC198
Memantine	Human Plasma	0.5-50 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Madison	Active	P786.01
	Human Plasma	0.5-50 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Madison	Active	P786.01
Metformin	Human Plasma	2-2,000 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Richmond	Active	LCMS153
	Human Plasma	2-1,000 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Richmond	Active	LCMS153.5
Metformin	Human Urine	10-10,000 µg/mL	HPLC/UV	0.05 mL	None	-20	Richmond	Active	LC251.1
	Dog Plasma	1-1,000 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Richmond	Active	LCMSB153.4

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Methadone	Human Plasma	0.1-50 ng/mL	LC/MS/MS	0.2 mL	K ₃ EDTA	-20	Richmond	Active	LCMS323
	Human Urine	25-1,000 ng/mL	HPLC/UV	1.0 mL	None	-20	Richmond	Inactive	LC238.2
Methadone Enantiomers	Human Plasma	5-1,000 ng/mL 5-1,000 ng/mL	LC/MS/MS	0.05 mL	K ₃ EDTA	-20	Richmond	Active	LCMSC323.1
Methotrexate (7-Hydroxymethotrexate)	Human Plasma	5-500 ng/mL 5-500 ng/mL	HPLC/UV	1.0 mL	Sodium Heparin	-20	Madison	Active	P461.02
	Human Urine	0.1-10 ug/mL 0.1-10 ug/mL	HPLC/UV	0.05 mL	None	-20	Madison	Active	P683.02
	Human Plasma	1.0-1,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Richmond	Active	LCMSC380.1
	Human Urine	0.05-25 ug/mL	LC/MS/MS	0.05 mL	None	-20	Richmond	Active	LCMSC380.2
	Human Plasma	1.0-1,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Richmond	Active	LCMS380
Methylphenidate	Human Plasma	0.25-25 ng/mL	LC/MS/MS	0.10 mL	Sodium Heparin	-20	Richmond	Inactive	LCMSB187
	Human Plasma	0.1-100 ng/mL	LC/MS/MS	0.05 mL	K ₃ EDTA	-70	Richmond	Active	LCMS230.1
Methylphenidate (Ritalinic Acid)	Beagle Plasma	0.10-100 ng/mL 2.0-2,000 ng/mL	LC/MS/MS	0.05 mL	K ₃ EDTA	-20	Richmond	Active	LCMS230
Methylphenidate (Ritalinic Acid)	Human Plasma	0.1-50 ng/mL 2.0-1,000 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P689.02

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Methylphenidate Enantiomers (Ritalinic Acid Enantiomers)	Human Plasma	0.05-50 ng/mL (D) 0.01-10 ng/mL (L) 1.0-1,000 ng/ml (D) 1.0-1,000 ng/ml (L)	LC/MS/MS	0.2 mL	Sodium Heparin	-20	Madison	Active	P779.00
Methyltestosterone	Human Plasma	5- 25,000 pg/mL	LC/MS/MS	0.5 mL	K ₃ EDTA	-20	Richmond	Active	LCMSC349
Metoprolol	Human Plasma	5-1,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Madison	Active	P940.00
Metronidazole (1-Hydroxymetronidazole)	Human Plasma	10-1,000 ng/mL 10-1,000 ng/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Madison	Active	P686.00
Mevalonic Acid	Human Plasma	0.5-20 ng/mL	LC/MS/MS	0.5 mL	K ₃ EDTA	-20	Madison	Active	P758.02
	Human Urine	25-1,000 ng/mL	LC/MS/MS	0.25 mL	None	-20	Madison	Active	P759.00
Midazolam (1-Hydroxy midazolam) (1-HydroxyMidazolam, Total**)	Human Plasma	0.1-100 ng/mL 0.1-100 ng/mL	LC/MS/MS	0.3 mL	Sodium Heparin	-20	Richmond	Active	LCMS97
	Human Urine	1-1,000 ng/mL	LC/MS/MS	0.2 mL	None	-20	Richmond	Inactive	LCMS228
Midazolam (1-Hydroxy midazolam) (4-Hydroxy midazolam)	Human Plasma	0.1-100 ng/mL 0.1-100 ng/mL 0.1-100 ng/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-20	Madison	Active	P1024.00
Midazolam (1-HydroxyMidazolam)	Human Urine	0.05-50 ng/mL 1-1,000 ng/mL	LC/MS/MS	0.2 mL	None	-20	Madison	Active	P678.04
Midazolam (1-HydroxyMidazolam)	Human Urine	0.05-50 ng/mL 2-2,000 ng/mL	LC/MS/MS	0.2 mL	None	-20	Madison	Active	P870.01
Minocycline	Human Plasma	0.02-10 µg/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Richmond	Active	LC63

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
	Human Serum	0.02-10 µg/mL	HPLC/UV	0.5 mL	None	-20	Richmond	Active	LC63.1
	Human Saliva	0.02-10 µg/mL	HPLC/UV	0.5 mL	None	-20	Richmond	Active	LC63.5
	Human Crevicular Fluid	10-2,000 ng/µL	HPLC/UV	1.0 µL	None	-20	Richmond	Active	LC360
Mirtazapine	Human Plasma	0.05-100 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Richmond	Active	LCMS145
Mometasone Furoate	Human Plasma	0.25-25 pg/mL	LC/MS/MS	1.0 mL	K ₃ EDTA	-20	Richmond	Active	LCMSD345.1
	Dog Plasma	10-1,000 pg/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Richmond	Active	LCMSD345.4
	Rat Plasma	10-1,000 pg/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Richmond	Active	LCMSD345.3
Montelukast	Human Plasma	5-1,000 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-80	Richmond	Active	LCMSC385
Morphine (Morphine-3-β-glucuronide) (Morphine-6-β-glucuronide)	Human Plasma	0.5-50 ng/mL 10-1,000 ng/mL 2-200 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Richmond	Active	LCMS 454
Morphine (Morphine-3-β-glucuronide) (Morphine-6-β-glucuronide) Codeine	Human Plasma	0.2-20 ng/mL 2-2,00 ng/mL 0.5-50 ng/mL 1-100 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Richmond	Active	LCMSC396
Morphine & Codeine & Guafenesin	Human Plasma	0.2-20 ng/mL 0.1-100 ng/mL 2-2,000 ng/mL	LC/MS/MS	0.3 mL	Sodium Heparin	-20	Madison	Active	P919.02
Moxifloxacin	Human Plasma	25-5,000 ng/mL	LC/MS/MS	0.05 mL	K ₃ EDTA	-20	Richmond	Active	LCMSB276

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Naloxone & Buprenorphine (Norbuprenorphine)	Human Plasma	5-500 pg/mL 50-10,000 pg/mL 50-10,000 pg/mL	LC/MS/MS	0.5 mL	K ₃ EDTA	-20	Madison	Active	P942.01
Naloxone, Total	Human Plasma	25-10,000 pg/mL	LC/MS/MS	0.2 mL	K ₃ EDTA	-20	Madison	Active	P943.00
Naproxen	Human Plasma	0.1-100 µg/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Richmond	Active	LCMSB486
Navelbine	Human Plasma	2-500 ng/mL	HPLC/FL	0.5 mL	Sodium Heparin	-70	Richmond	Inactive	LC253.1
Nevirapine	Human Plasma	25-10,000 ng/mL	LC/MS/MS	0.05 mL	K ₃ EDTA	-20	Richmond	Active	LCMS250
Niacin <i>aka Vitamin B3, or nicotinic acid</i> (Nicotinuric acid) <i>aka NUA</i> (1-Methyl-Nicotinamide) <i>aka NMA</i> (N-Methyl-2-Pyridone-5-carboxamide) <i>aka 2PY</i>	Human Plasma	5-2,000 ng/mL 5-2,000 ng/mL	LC/MS/MS	0.25mL	Sodium Heparin	-70	Richmond	Active	LCMSB264
	Human Urine	0.02 -20 µg/mL 0.2 - 200 µg/mL	LC/MS/MS	0.1 mL	None	-20	Richmond	Active	LCMSB264.1
	Human Urine	0.5-250 µg/mL 2.5-250 µg/mL	LC/MS/MS	0.05 mL	None	-20	Richmond	Active	LCMS274.1
	Human Plasma	0.1-10 µg/mL 0.2- 20 µg/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Richmond	Active	LCMSB274
(1-Methyl-Nicotinamide) <i>aka NMA</i> (N-Methyl-2-Pyridone-5-carboxamide) <i>aka 2PY</i>	Human Plasma	0.1-10 µg/mL 0.2- 20 µg/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Richmond	Active	LCMSB274.2
Nicotine (Cotinine)	Human Plasma	0.5-100 ng/mL 2.5-500 ng/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Madison	Active	P538.02
Nilotinib	Human Plasma	2-2,000 ng/mL	LC/MS/MS	0.05 mL	K₂EDTA	-20	Madison	Active	P1011

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Nizatidine (N-Desmethylnizatidine)	Human Plasma	5-5,000 ng/mL 5-5,000 ng/mL	HPLC/UV	1.0 mL	Sodium Heparin	-20	Richmond	Active	LC264
Norelgestromin <i>aka 17-desacetyl norgestimate</i>	Human Plasma	0.02-10 ng/mL	LC/MS/MS	0.5 mL	Potassium Oxalate/Sodium Fluoride	-70	Richmond	Active	LCMSC372
Norethindrone & Ethinyl Estradiol	Human Plasma	50-25,000 pg/mL 2-500 pg/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Richmond	Active	LCMSC255
Norgestrel & Ethinyl Estradiol	Human Plasma	50-25,000 pg/mL 2-500 pg/mL	LC/MS/MS	0.5 mL	Potassium Oxalate/Sodium Fluoride	-70	Richmond	Active	LCMSC256
Octreotide	Human Plasma	0.025-25 ng/mL	LC/MS/MS	0.2 mL	K ₃ EDTA	-20	Richmond	Active	LCMSC296
	Monkey Plasma	5.0-250 ng/mL	LC/MS/MS	0.2 mL	K ₃ EDTA	-70	Richmond	Active	LCMSC296.2
	Monkey Plasma	0.1-25 ng/mL	LC/MS/MS	0.2 mL	K ₃ EDTA	-70	Richmond	Active	LCMSC296.1
	Rabbit Plasma	1-100 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-70	Richmond	Active	LCMSC296.3
Olanzapine (Desmethyloanzapine)	Human Plasma	0.05-50 ng/mL 0.1-100 ng/mL	LC/MS/MS	0.3 mL	Sodium Heparin	-20	Richmond	Active	LCMS84
Omeprazole	Human Plasma	1-1,000 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Richmond	Active	LCMSB131
	Equine Plasma	1-1,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Madison	Active	P1025.00
Omeprazole (5-Hydroxyomeprazole) (Omeprazole sulphone)	Human Plasma	1-1,000 ng/mL 1-1,000 ng/mL 1-1,000 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P749.04

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Ondansetron	Human Plasma	0.5-250 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Richmond	Active	LCMS165
	Dog Plasma	0.5-250 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Richmond	Active	LCMS165.1
Oxaprozin	Human Plasma	0.5-500 µg/mL	HPLC/UV	0.1 mL	Sodium Heparin	-20	Richmond	Inactive	LC243
Oxcarbazepine (10-Hydroxycarbamazepine (MHD)) (10,11-Dihydroxycarbamazepine (DHD))	Human Plasma	0.1-5 µg/mL 0.1-5 µg/mL 0.5-25 µg/mL	HPLC/UV	0.3 mL	Sodium Heparin	-70	Madison	Active	P743.00
	Human Plasma	0.1-5 µg/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-70	Madison	Active	P850.00
(Monohydroxyoxcarbazepine <i>aka</i> MHD)	Human Plasma	1-50 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-70	Madison	Active	P886.00
Oxybutynin Enantiomers (Desethyloxybutynin enantiomers)	Human Plasma	0.5-200 ng/mL 0.5-200 ng/mL	LC/MS/MS	0.2 mL	K ₂ EDTA	-70	Richmond	Active	LCMS 411
Oxycodone	Human Plasma	0.1-100 ng/mL	LC/MS/MS	0.5 mL	K ₃ EDTA	-20	Madison	Active	P649.01
Oxymorphone	Human Plasma	10-1,000 pg/mL	LC/MS/MS	0.5 mL	K ₃ EDTA	-20	Madison	Active	P820.01
Oxymorphone (6-OH-oxymorphone)	Human Plasma	0.1-10 ng/mL 0.1-10 ng/mL	LC/MS/MS	0.25 mL	K₂EDTA	-20	Madison	Active	P930
Paclitaxel (Taxol ®)	Human Plasma	5-1,000 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-70	Richmond	Active	LCMS163
	Human Plasma	0.1-100 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Richmond	Active	LCMS163.2
Paroxetine	Human Plasma	0.25-50 ng/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-20	Richmond	Active	LCMS161

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
	Human Plasma	0.25-50 ng/mL	LC/MS/MS	0.2 mL	Sodium Heparin K ₃ EDTA	-20	Madison	Active	P776.01
Pemoline	Human Plasma	25-5,000 ng/mL	HPLC/UV	0.2 mL	Sodium Heparin	-20	Richmond	Inactive	LC258
Pentoxifylline (Hydroxypentifylline) (Carboxypropylpentifylline)	Human Plasma	5-1,000 ng/mL 10-2,000 ng/mL 10-2,000 ng/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Madison	Active	P348.01
Phenethylamine	Human Urine	1.13-501 ng/mL	LC/MS/MS	0.2 mL	None	-20	Madison	Active	P655.00
Phenobarbital See Cocktail Assay "Epilepsy"	Human Plasma	0.1-50 µg/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-70	Madison	Active	P837.00
Phenylbutazone	Equine Plasma	0.1-40 µg/mL	HPLC/UV	0.25 mL	Sodium Heparin	-20	Madison	Active	P554.04
Phenylephrine, Total**	Human Plasma	5.0-1,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-70	Madison	Active	P806.01
	Human Plasma	1.0-1,000 ng/mL	LC/MS/MS	0.05 mL	Lithium Heparin	-70	Richmond	Active	LCMS257
	Human Urine	100-5,000 ng/mL	LC/MS/MS	0.05 mL	None	-70	Madison	Active	P958.01
Phenylephrine (Phenylephrine Glucuronide)	Human Urine	2.0-200 ng/mL 2.0-200 ng/mL	LC/MS/MS	0.2 mL	None	-70	Madison	Active	P956.00
(3-Hydroxymandelic acid)	Human Urine	100-5,000 ng/mL	LC/MS/MS	0.05 mL	None	-70	Madison	Active	P957.00
Phenylephrine	Human Plasma	10-2,500 pg/mL	LC/MS/MS	0.25 mL	Lithium Heparin	-70	Richmond	Active	LCMSC392.1
	Human Plasma	10-2,500 pg/mL	LC/MS/MS	0.25 mL	K ₃ EDTA	-70	Madison	Active	P898.01
Phenylpropanolamine	Human Plasma	5-1,000 ng/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Richmond	Inactive	LC181

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
	Human Plasma	5-500 ng/mL	HPLC/UV	1.0 mL	EDTA	-20	Madison	Inactive	P002.05
Phenytoin	Human Plasma	20-5,000 ng/mL	LC/MS/MS	0.25 mL	K ₂ EDTA	-20	Richmond	Active	LCMSB418
Phenytoin Phenytoin, Total **	Human Plasma	0.05-15 ug/mL 0.1- 30 ug/mL	LC/MS/MS	0.05 mL 0.025 mL	K ₂ EDTA	-20	Madison	Active	P639.00
Phenytoin, Free (unbound) (5-Hydroxyphenytoin)	Human Plasma Ultrafiltrate	0.05-15 µg/mL 0.05-15 µg/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Madison	Active	P742.03
Phenytoin, Total** (5-Hydroxyphenytoin)	Human Plasma	0.05-15 µg/mL 0.1- 30 µg/mL	LC/MS/MS	0.025 mL	Sodium Heparin	-20	Madison	Active	P742.03
(5-Hydroxyphenytoin)	Human Plasma	0.1-20 µg/mL	HPLC/UV	1.0 mL	Sodium Heparin	-20	Madison	Active	P281.00
Phenytoin (5-Hydroxyphenytoin)	Human Urine	0.1-40 µg/mL 0.1-40 µg/mL	HPLC/UV	0.5 mL	None	-20	Madison	Active	P642.01
(5-Hydroxyphenytoin), Total**	Human Urine	1-200 µg/mL	HPLC/UV	0.05 mL	None	-20	Madison	Active	P643.00
Pravastatin	Human Plasma	0.5-250 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Richmond	Active	LCMS47
Praziquantel	Dog Plasma	10-4,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Madison	Active	P989.00
	Feline Plasma	10-4,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Madison	Active	P831.01
Prednisone (Prednisolone)	Human Plasma	0.5-1,000 ng/mL 0.5-1,000 ng/mL	LC/MS/MS	0.3 mL	K ₂ EDTA	-20	Madison	Active	P955.00
	Rat Plasma	0.25-250 ng/mL 0.25-250 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-70	Madison	Active	P984.00

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Pregabalin	Human Plasma	20-20,000 ng/mL	LC/MS/MS	0.5 mL	K ₂ EDTA	-20	Richmond	Active	LCMSC487
Procainamide (N-Acetylprocainamide)	Human Plasma	0.025-10 µg/mL 0.025-10 µg/mL	HPLC/UV	0.25 mL	Sodium Heparin	-20	Richmond	Inactive	LC306
Progesterone	Human Plasma	0.05-50 ng/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-70	Richmond	Active	LCMSC431
	Rabbit Plasma	0.5-50 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-70	Richmond	Active	LCMSC431.1
Proguanil	Human Plasma	0.5-500 ng/mL	LC/MS/MS	0.05 mL	K₂EDTA	-20	Richmond	Active	LCMS530
Propranolol (4-Hydroxypropranolol)	Human Plasma	0.5-250 ng/mL 0.5-250 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-70	Richmond	Active	LCMSB191
	Human Urine	1-250 ng/mL 1-250 ng/mL	LC/MS/MS	0.25 mL	None	-70	Richmond	Active	LCMSB191.1
Propranolol, Total** (4-Hydroxypropranolol)	Human Plasma	2-600 ng/mL 2-600 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-70	Richmond	Active	LCMSB281.2
Pseudoephedrine	Human Plasma	2.5-500 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Richmond	Active	LCMSB201
Pseudoephedrine & Dextromethorphan	Human Plasma	0.8-400 ng/mL 10-5,000 pg/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Madison	Active	P617.02
Quetiapine	Human Plasma	1-1000 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Richmond	Active	LCMSB430
Quinidine	Human Plasma	25-5,000 ng/mL	HPLC/FL	0.5 mL	Sodium Heparin	-20	Richmond	Inactive	LC330.1
Raltegravir See Cocktail Assay "Antiviral"	Human Plasma	10-10,000 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Madison	Active	P1002.00

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Ramipril (Ramiprilat)	Human Plasma	0.05-50 ng/mL 0.05-50 ng/mL	LC/MS/MS	0.25 mL	K ₃ EDTA	-70	Richmond	Active	LCMS311
Ranitidine	Human Plasma	2.5-2,500 ng/mL	LC/MS/MS	0.05 mL	K ₃ EDTA	-20	Madison	Active	P873.01
Ranolazine	Human Plasma	2-2,000 ng/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Madison	Active	P934.01
Ranolazine (RS-94827 Metabolite)	Human Plasma	2-2,000 ng/mL 2-2,000 ng/mL	LC/MS/MS	0.25 mL	K₂EDTA	-20	Madison	Active	P1015
Repaglinide	Human Plasma	0.2-200 ng/mL	LC/MS/MS	0.1 mL	K ₂ EDTA	-20	Richmond	Active	LCMS460
Ribavirin	Human Plasma	5-5,000 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Richmond	Active	LCMS510
Rifabutin (25-O-Desacetylriofabutin)	Human Plasma	2-800 ng/mL 2-800 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Madison	Active	P587.06
Rifampin	Human Plasma	0.1-10 µg/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Madison	Active	P941.02
Risperidone (9-Hydroxyrisperidone)	Human Urine	1-300 ng/mL 1-300 ng/mL	LC/MS/MS	0.2 mL	None	-20	Madison	Active	P362.01
	Human Plasma	0.1-60 ng/mL 0.1-60 ng/mL	LC/MS/MS	0.1 mL	K₂EDTA	-20	Madison	Active	P1019
	Human Serum	0.1-30 ng/mL 0.1-30 ng/mL	LC/MS/MS	1.0 mL	None	-20	Madison	Active	P361.00
Ritonavir	Human Plasma	10-10,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-70	Richmond	Active	LCMS210.2

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
See Cocktail Assay "Antiviral"	Human Plasma	10-10,000 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Madison	Active	P1002.00
Ritonavir & Atazanavir	Human Plasma	10-10,000 ng/mL	LC/MS/MS	0.05 mL	K₃EDTA	-20	Madison	Active	P1012
Rivaroxaban	Human Plasma	0.5-500 ng/mL	LC/MS/MS	0.25 mL	K ₂ EDTA	-20	Madison	Active	P1014.00
Rizatriptan	Human Plasma	0.2-200 ng/mL	LC/MS/MS	0.05 mL	K ₃ EDTA	-20	Madison	Active	P845.00
Ropinirole	Human Plasma	10-5,000 pg/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-20	Madison	Active	P736.01
Rosiglitazone (N-Desmethylrosiglitazone)	Human Plasma	1-500 ng/mL 1-500 ng/mL	LC/MS/MS	0.05 mL	K ₃ EDTA	-20	Richmond	Active	LCMS322.1
Rosiglitazone	Human Plasma	2-1,000 ng/mL	LC/MS/MS	0.05 mL	K ₃ EDTA	-20	Richmond	Active	LCMS322
Rosiglitazone	Rat Plasma	2-1,000 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Richmond	Active	LCMS322.2
Rosuvastatin	Human Plasma	0.1-100 ng/mL	LC/MS/MS	0.2 mL	K ₂ EDTA	-70	Richmond	Active	LCMSC393
Rosuvastatin (N-Desmethylrosuvastatin)	Human Plasma	0.1-100 ng/mL 0.05-50 ng/mL	LC/MS/MS	0.25 mL	K ₃ EDTA	-70	Madison	Active	P922.03
(Salicylic Acid)	Human Plasma	1-50 µg/mL	HPLC/UV	0.2 mL	Lithium Heparin	-20	Madison	Active	P383.00
	Dog Plasma	1-50 µg/mL	HPLC/UV	0.2 mL	Sodium Heparin	-70	Richmond	Inactive	LC261.1
Acetylsalicylic Acid (Salicylic Acid)	Human Plasma	0.02-10 µg/mL 0.1-50 µg/mL	LC/MS/MS	0.05 mL	Potassium Oxalate/Sodium Fluoride	-70	Richmond	Active	LCMSC425

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Salmeterol	Human Plasma	1-100 pg/mL	LC/MS/MS	1.0 mL	Potassium Oxalate/Sodium Fluoride	-20	Madison	Active	P798.00
Salmeterol & Fluticasone	Human Plasma	1-200 pg/mL 1-200 pg/mL	LC/MS/MS	1.0 mL	Potassium Oxalate/Sodium Fluoride	-20	Madison	Active	P988
Selegiline (N-Desmethylselegiline) (Amphetamine) (Methamphetamine) (Phenethylamine)	Human Plasma	0.01-5 ng/mL 0.04-20 ng/mL 0.05-25 ng/mL 0.05-25 ng/mL 0.01-5 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P985.00
Sertraline	Human Plasma	0.2-80 ng/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Madison	Active	P570.01
Sertraline (Desmethylsertraline)	Human Plasma	0.5-200 ng/mL 0.5-200 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Richmond	Active	LCMS98.1
Sildenafil (Desmethylsildenafil)	Human Plasma	1-1,000 ng/mL 1-1,000 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin & K ₃ EDTA	-70	Richmond	Active	LCMS231
Simvastatin (Simvastatin Acid)	Human Plasma	0.1-50 ng/mL 0.1-50 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-70	Richmond	Active	LCMSC367
	Human Plasma	0.1-50 ng/mL 0.1-50 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-70	Madison	Active	P980.00
Solifenacin	Human Plasma	0.1-25 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-20	Madison	Active	P950.00
Sotalol	Human Plasma	10-2,000 ng/mL	HPLC/FL	0.5 mL	EDTA	-20	Madison	Active	P403.01
Sparfloxacin	Human Plasma	0.025-20 µg/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Richmond	Active	LC332

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Sufentanil	Human Plasma	1.0-250 pg/mL	LC/MS/MS	0.5 mL	K ₂ EDTA	-70	Richmond	Active	LCMSC239
	Human Urine	5.0-1,000 pg/mL	LC/MS/MS	0.25 mL	None	-70	Richmond	Active	LCMSC239.1
Sumatriptan	Human Plasma	0.2-100 ng/mL	LC/MS/MS	0.25 mL	K ₃ EDTA	-20	Richmond	Active	LCMS174
	Pig Plasma	0.2-100 ng/mL	LC/MS/MS	0.25 mL	K ₂ EDTA	-20	Richmond	Active	LCMS174.3
Sunitinib (N-Desethylsunitinib)	Human Plasma	0.1-50 ng/mL 0.1-50 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Richmond	Active	LCMSC457
T3	Human Serum	0.25-8 ng/mL	RIA	0.2 mL	None	-20	Richmond	Active	ICD86
T4	Human Serum	1-32 µg/dL	RIA	0.05 mL	None	-20	Richmond	Active	ICD87
Tacrolimus	Human Whole Blood	0.25-100 ng/mL	LC/MS/MS	0.25 mL	K ₃ EDTA	-20	Richmond	Active	LCMS356
Tamsulosin	Human Plasma	0.05-25 ng/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Madison	Active	P682.02
Tamsulosin	Dog Plasma	0.05-25 ng/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Madison	Active	P750.01
	Pig Plasma	0.05-25 ng/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Madison	Active	P969.00
Telmisartan	Human Plasma	2-1,000 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Richmond	Active	LCMSC408
Tenofovir & Emtricitabine	Human Plasma	5-1,000 ng/mL 20-4,000 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Richmond	Active	LCMSC 468
	Human Plasma	5-1,000 ng/mL 20-4,000 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Madison	Active	P947.00

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Terfenadine (Fexofenadine)	Human Plasma	0.05-50 ng/mL 0.5-500 ng/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Madison	Active	P473.01
Terazosin	Human Plasma	0.5-200 ng/mL	HPLC/FL	0.3 mL	Sodium Heparin	-20	Richmond	Inactive	LC270
Terconazole	Balb/c Mouse Plasma	1-100 ng/mL	LC/MS/MS	0.025 mL	K ₃ EDTA	-20	Richmond	Active	LCMSC217.1
Testosterone (Dihydrotestosterone)	Human Serum	50–10,000 pg/mL 50 -10,000 pg/mL	LC/MS/MS	0.15 mL	None	-20	Richmond	Active	LCMSC260.6
Testosterone, Free (unbound)	Human Serum Ultrafiltrate	1-200 pg/mL	LC/MS/MS	0.2 mL	None	-20	Richmond	Active	LCMSC247
Theophylline	Human Plasma	0.05-25 µg/mL	HPLC/UV	0.5 mL	Sodium Heparin	-20	Richmond	Inactive	LC9
	Human Serum	0.05-25 µg/mL	HPLC/UV	0.5 mL	None	-20	Richmond	Inactive	LC9.1
TNF-α	Human Serum	15.6-1,000 pg/mL	ELISA	0.4 mL	None	-20	Richmond	Active	ICD134
Tobramycin	Human Plasma	10-400 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P790.00
Tobramycin	Human Urine	100-20,000 ng/mL	LC/MS/MS	0.05 mL	None	-20	Madison	Active	P769.01
Tolbutamide	Human Plasma	0.1-100 µg/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Richmond	Active	LCMSB226
Tolbutamide (Carboxytolbutamide) (4-Hydroxytolbutamide)	Human Urine	5-1,000 ng/mL 0.6-300 µg/mL 0.3-150 µg/mL	LC/MS/MS	0.02 mL	None	-20	Madison	Active	P738.01
	Human Plasma	0.1-100 µg/mL 5-5,000 ng/mL 2.5-2,500 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P737.02

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Topiramate	Human Plasma	10-10,000 ng/mL	LC/MS/MS	0.1mL	K ₂ EDTA	-20	Richmond	Active	LCMS470
Topiramate (See Cocktail Assay "Epilepsy")	Human Plasma	0.1-50 µg/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-70	Madison	Active	P837.00
Topotecan, Total	Human Plasma	0.1-25 ng/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-70	Richmond	Active	LCMSD507
Topotecan Lactone	Human Plasma	0.02-5 ng/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-70	Richmond	Active	LCMSD504
Trastuzumab aka Herceptin	Human Serum	10-200 µg/mL	ELISA	0.2 mL	None	-70	Richmond	Active	ICD178
	Human Plasma	10-100 µg/mL	ELISA	0.1 mL	EDTA	-20	Richmond	Active	ICD178.1
Triamcinolone	Human Plasma	0.01-2.5 ng/mL	LC/MS/MS	0.25 mL	K ₃ EDTA	-70	Richmond	Active	LCMSC315
	Human Plasma	1-1,000 ng/mL	LC/MS/MS	0.1 mL	K ₃ EDTA	-70	Richmond	Active	LCMSC315.1
Triamterene (Hydroxytriamterene Sulfate)	Human Plasma	0.5-100 ng/mL 5-1,000 ng/mL	HPLC/FL	0.5 mL	Sodium Heparin	-20	Richmond	Inactive	LC109
Triamterene (Hydroxytriamterene Sulfate)	Human Urine	0.10-5 µg/mL 1-50 µg/mL	HPLC/FL	0.1 mL	None	-20	Richmond	Inactive	LC109.1
TSH	Human Serum	0.250-64 µIU/mL	RIA (IRMA)	0.4 mL	None	-20	Richmond	Inactive	ICD179
Valganciclovir (Ganciclovir)	Human Plasma	1-500 ng/mL 10-5,000 ng/mL	LC/MS/MS	0.05 mL	Sodium Citrate	-70	Richmond	Active	LCMSC410
Valproic Acid	Human Plasma	2-100 µg/mL	LC/MS/MS	0.05 mL	Sodium Heparin	-20	Madison	Active	P885.00
Venlafaxine (O-Desmethylvenlafaxine)	Human Plasma	2-200 ng/mL 2-200 ng/mL	LC/MS/MS	0.5 mL	Sodium Heparin	-20	Richmond	Active	LCMS531

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list



LIST OF VALIDATED METHODS JUNE 2010

Compound (Metabolite)	Matrix	Calibration Range	Method	Sample Volume	Anti-Coagulant	Storage Temp ° C	Lab Site	Method Status*	Method I.D.
Verapamil (Norverapamil)	Human Plasma	1-500 ng/mL 1-500 ng/mL	HPLC/FL	1.0 mL	Sodium Heparin	-20	Madison	Inactive	P072.02
Verapamil Enantiomers (Norverapamil Enantiomers)	Human Plasma	0.5-250 ng/mL 0.5-250 ng/mL	LC/MS/MS	0.4 mL	Sodium Heparin	-20	Madison	Active	P671.00
Voriconazole (Voriconazole-N-Oxide)	Human Plasma	10-5,000 ng/mL 10-5,000 ng/mL	LC/MS/MS	0.1 mL	Sodium Heparin	-20	Madison	Active	P968.00
Warfarin Enantiomers, (unbound)	Human Plasma Ultrafiltrate	0.5-25 ng/mL 0.5-25 ng/mL	LC/MS/MS	0.05 mL	EDTA	-20	Madison	Active	P921.00
Warfarin Enantiomers	Human Serum	5-1,000 ng/mL 5-1,000 ng/mL	LC/MS/MS	0.5 mL	None	-20	Madison	Inactive	P381.01
	Human Plasma	5-1,000 ng/mL 5-1,500 ng/mL	LC/MS/MS	0.25 mL	Sodium Heparin	-20	Richmond	Active	LCMSB194
Warfarin Enantiomers	Human Plasma	5-1,500 ng/mL 5-1,500 ng/mL	LC/MS/MS	0.2 mL	Sodium Heparin	-20	Madison	Active	P807.01
S-warfarin (7-OH-S-warfarin)	Human Plasma	5-1,500 ng/mL 5-1,000 ng/mL	LC/MS/MS	0.2 mL	Sodium Citrate	-20	Madison	Active	P832.00
Zolpidem	Human Plasma	0.5-500 ng/mL	LC/MS/MS	0.05 mL	K ₂ EDTA	-20	Madison	Active	P844.01
Zonisamide	Human Plasma	10-5,000 ng/mL	LC/MS/MS	0.25 mL	K ₃ EDTA	-20	Richmond	Active	LCMS313
	Human Serum	10-5,000 ng/mL	LC/MS/MS	0.25 mL	None	-20	Richmond	Active	LCMS313.1

*Inactive methods and older methods may require a revalidation fee to bring them up-to-date with the current FDA Guidance

**Total = Unconjugated + Conjugated Drug.

Unless specifically noted, the free or unconjugated drug is measured.

Shaded entries are NEW to our list